

Unusual Two-stage Dynamics of the Spin-Lattice Polaron Formation

J. Kogoj^{1,1}, Z. Lenarčič¹, D. Golež¹, M. Mierzejewski^{2,3}, P. Prelovšek^{1, 2} and J. Bonča^{1, 2}

¹*J. Stefan Institute, 1000 Ljubljana, Slovenia;*

²*Faculty of Mathematics and Physics, University of Ljubljana, 1000 Ljubljana, Slovenia;*

³*Institute of Physics, University of Silesia, 40-007 Katowice, Poland;*

I will present a fundamental study of a Holstein polaron in one dimension and a single hole in the two dimensional t - J -Holstein model driven by a constant electric field. Taking fully into account quantum effects we follow the time-evolution of systems from their ground state as the electric field is switched on at $t = 0$, until they reach a steady state. In the Holstein polaron we observe: adiabatic regime with Bloch oscillations and zero net current and a dissipative regime with a finite current that is further divided on a linear $I - V$ region and a region with a negative differential resistance [1,2].

In the case you found this abstract tiresome, the scene on Fig.1 might convince you of joining the conference.

Figure 1: The idyllic scene of Krvavec and hotel "Raj". Disclaimer: This figure might not reflect the actual scene at the time of the conference.

References

- [1] L. Vidmar, J. Bonča, M. Mierzejewski, P. Prelovšek, and S.A. Trugman, *Phys. Rev. B* **83** 134301, (2011).
- [2] M. Mierzejewski, L. Vidmar, J. Bonča, and P. Prelovšek, *Phys. Rev. Lett.* **106** 196401, (2011).

¹jan.kogoj@ijs.si